

Driving Safety and Medicines

Did you know that some over-the-counter and prescription medicines can affect the way you drive?

Medicines can affect people in different ways.

Ask your doctor or pharmacist for more information.

What types of medicines can affect my driving?

Some examples include:

- Some cold remedies and allergy products (e.g. Codral Day and Night[®], Phenergan[®])
- Sleeping tablets and anxiety medicine (e.g. Antenex[®]/Valium[®], Stilnox[®])
- Pain and opioid medicines (e.g. Lyrica[®], Panadeine Forte[®], Endone[®], Tramal[®])
- Some antidepressant and antipsychotic medicines (e.g. Seroquel[®], Lexapro[®])
- Some medicines for epilepsy (e.g. Neurontin[®])
- Opioid treatment medicines methadone and buprenorphine (e.g. Suboxone[®], Subutex[®])
- Prescribed cannabis medicines

Drinking alcohol or taking illicit drugs when taking medicines, taking more than the prescribed dose, or taking a combination of these substances can seriously affect your ability to drive safely.



How can these medicines affect my driving?

Some medicines can cause:

- Confusion/poor concentration
- Slower reaction time
- Drowsiness/tiredness
- Dizziness/feeling faint
- Blurred or double vision
- Muscle weakness
- Anxiety and mood changes
- Shakiness/unsteadiness

These symptoms can make it unsafe for you to drive, cycle or use machinery and can put you and other road users at risk of an accident.

If you are taking any medicine and feel any of the symptoms listed above or are unsure about driving

DON'T DRIVE



